


Amendments to the Claims:

This listing of claims will replace all prior versions, and listings of claims in the application:

Listing of Claims:

Please amend the claims as follows:

1. (Original) Process for preparing a Palladium(0) compound, comprising reaction of a palladium compound with one or more compounds of the general formula I in the presence of a base:



in which:

each A is independently a CR^7R^8 -radical where one of the A radicals may be oxygen, sulphur, an NR^9 group or an $\text{SiR}^{10}\text{R}^{11}$ group, or where the A radicals may be a constituent of a 5- to 20-membered ring system,


x is an integer from 2 to 4, and

each R^1 to R^{11} is independently selected from R, OR, halogen, CN, NO_2 , NR_2 , C(O)R , C(O)OR , OC(O)R , CONR_2 , NHCO_2R , NHCOR , $\text{CH=CH-CO}_2\text{R}$, Si(R)_3 , Si(OR)_3 , SiR(OR)_2 , $\text{SiR}_2(\text{OR})$, SO_3R , SO_2R , SOR , SR , PR_2 , POR_2 , PO_3H , PO(OR)_2 , in which R is a hydrogen atom, a substituted or unsubstituted C_{1-10} -alkyl radical, a substituted or

unsubstituted, mono- or polyunsaturated C₁₋₁₀-alkenyl radical, or a substituted or unsubstituted, optionally heteroatom-containing C₅₋₁₀-aryl radical, and the substituents on the alkyl radical or the alkenyl radical are selected from halogen, O-C₁₋₁₀-alkyl, phenyl, O-phenyl, OH, NH₂ and halogenated C₁₋₁₀-alkyl, and the substituents on the aryl radical are selected from halogen, C₁₋₁₀-alkyl, O-C₁₋₁₀-alkyl, phenyl, O-phenyl, OH, NH₂ and halogenated C₁₋₁₀-alkyl, where R² and R³ and/or R⁵ and R⁶ together with the carbon atoms bonded thereto may be a constituent of a 5- to 7-membered, optionally heteroatom-containing ring.

2. (Original) Process according to Claim 1, wherein x is 3.
3. (Previously Presented) Process according to Claim 1, wherein R¹ to R⁶ are each hydrogen atoms.
4. (Previously Presented) Process according to Claim 1, wherein -(A)_x- is a group of the formula -CH₂-X-CH₂- and -X- is selected from -O-, -S-, -SiR₂-, -NR- and -NC(O)R, and R is hydrogen, a C₁₋₄-alkyl radical or a halogenated C₁₋₄-alkyl.
5. (Previously Presented) Process according to Claim 1, wherein the compound of the general formula I is selected from the group consisting of 1,5-hexadiene, 1,6-heptadiene and 1,7-octadiene.
6. (Previously Presented) Process according to Claim 1, wherein the compound of the general formula I is selected from the group consisting of diallyl ether, diallylamine, diallylmethylamine, diallylethylamine, N-acetyldiallylamine, diallyl sulphide, diallylsilane, diallyldimethylsilane, difurfuryl ether, difurfurylamine, bis(thiophen-2-ylmethyl)amine, difurfuryl sulphide and 1,3-divinylbenzene.

7. (Original) Process for preparing a Palladium(0) compound, comprising reaction of a palladium compound with one or more compounds of the general formula II in the presence of a base:



in which:


n is an integer from 3 to 20,

each R^{13} to R^{15} is independently selected from R, OR, halogen, CN, NO_2 , NR_2 , $\text{C}(\text{O})\text{R}$, $\text{C}(\text{O})\text{OR}$, $\text{OC}(\text{O})\text{R}$, CONR_2 , NHCO_2R , NHCOR , $\text{CH}=\text{CH}-\text{CO}_2\text{R}$, $\text{Si}(\text{R})_3$, $\text{Si}(\text{OR})_3$, $\text{SiR}(\text{OR})_2$, $\text{SiR}_2(\text{OR})$, SO_3R , SO_2R , SOR , SR , PR_2 , POR_2 , PO_3H , $\text{PO}(\text{OR})_2$, in which R is hydrogen, a substituted or unsubstituted C_{1-10} -alkyl radical, a substituted or unsubstituted, mono- or polyunsaturated C_{1-10} -alkenyl radical, or a substituted or unsubstituted, optionally heteroatom-containing C_{5-10} -aryl radical, and the substituents on the alkyl radical or the alkenyl radical are selected from halogen, O- C_{1-10} -alkyl, phenyl, O-phenyl, OH, NH_2 and halogenated C_{1-10} -alkyl, and the substituents on the aryl radical are selected from halogen, C_{1-10} -alkyl, O- C_{1-10} -alkyl, phenyl, O-phenyl, OH, NH_2 and halogenated C_{1-10} -alkyl, where R^{13} and R^{14} together with the carbon atoms bonded thereto may be a constituent of a 5- to 7-membered, optionally heteroatom-containing ring, and

each R^{12} is independently selected from hydrogen, a hydroxyl group, a substituted or unsubstituted C_{1-10} -alkyl radical, an -O- C_{1-10} -alkyl radical (where the alkyl radical may be substituted or unsubstituted), a substituted or unsubstituted, mono- or polyunsaturated

C₁₋₁₀-alkenyl radical or a substituted or unsubstituted, optionally heteroatom-containing C₅₋₁₀-aryl radical, where the substituents are as defined for R¹³ and R¹⁵.

8. (Original) Process according to Claim 7, wherein n is an integer from 3 to 6 and each R¹² is independently a C₁₋₄-alkyl radical or a halogenated C₁₋₄-alkyl radical.
9. (Original) Process for preparing a Palladium(0) compound, comprising reaction of a palladium compound with one or more compounds of the general formula III in the presence of a base:



in which

v and w are each independently 0 or an integer of from 1 to 1000 and v+w is from 0 to 1000,


each R¹⁶ is independently selected from hydrogen, a hydroxyl group, a substituted or unsubstituted C₁₋₁₀-alkyl radical, an -O-C₁₋₁₀-alkyl radical (where the alkyl radical may be substituted or unsubstituted), a substituted or unsubstituted, mono- or polyunsaturated C₁₋₁₀-alkenyl radical or a substituted or unsubstituted, optionally heteroatom-containing C₅₋₁₀-aryl radical, where the substituents are as defined for R¹⁷ and R¹⁹,

each R¹⁷ to R¹⁹ is independently selected from R, OR, halogen, CN, NO₂, NR₂, C(O)R, C(O)OR, OC(O)R, CONR₂, NHCO₂R, NHCOR, CH=CH-CO₂R, Si(R)₃, Si(OR)₃, SiR(OR)₂, SiR₂(OR), SO₃R, SO₂R, SOR, SR, PR₂, POR₂, PO₃H, PO(OR)₂, in which R is a hydrogen atom, a substituted or unsubstituted C₁₋₁₀-alkyl radical, a substituted or unsubstituted, mono- or polyunsaturated C₁₋₁₀-alkenyl radical, or a substituted or unsubstituted, optionally heteroatom-containing C₅₋₁₀-aryl radical, and the substituents on the alkyl radical or the alkenyl radical are selected from halogen, O-C₁₋₁₀-alkyl, phenyl,

O-phenyl, OH, NH₂ and halogenated C₁₋₁₀-alkyl, and the substituents on the aryl radical are selected from halogen, C₁₋₁₀-alkyl, O-C₁₋₁₀-alkyl, phenyl, O-phenyl, OH, NH₂ and halogenated C₁₋₁₀-alkyl, where R¹⁷ and R¹⁹ together with the carbon atoms bonded thereto may be a constituent of a 5- to 7-membered, optionally heteroatom-containing ring,

each R²⁰ is independently selected from hydrogen, a hydroxyl group, a substituted or unsubstituted C₁₋₁₀-alkyl radical, an -O-C₁₋₁₀-alkyl radical (where the alkyl radical may be substituted or unsubstituted), a substituted or unsubstituted, mono- or polyunsaturated C₁₋₁₀-alkenyl radical, or a substituted or unsubstituted, optionally heteroatom-containing C₅₋₁₀-aryl radical, where the substituents are as defined for R¹⁷ and R¹⁹, and each Term is independently (R¹⁶)₂(CR¹⁷R¹⁸CR¹⁹)Si- or (R¹⁶)₃Si-.

10. (Original) Process according to Claim 9, wherein the compound of the general formula (III) has the general formula:



where R¹⁶ to R¹⁹, Term and v are each as defined in Claim 9.

11. (Previously Presented) Process according to Claim 7, wherein the compound of the general formula II or III is selected from divinylidisiloxane, 1,1,3,3-tetramethyl-1,3-divinylidisiloxane, 1,1,3,3-tetramethyl-1,3-dithien-2-ylidisiloxane, 1,1,3,3-tetramethoxy-1,3-divinylidisiloxane, 1,3-dimethyl-1,3-divinylidisiloxanediol, 1,3,5,7-tetravinyl-1,3,5,7-tetramethylcyclotetrasiloxane and 1,3,5-trimethyl-1,3,5-trivinylcyclotrisiloxane.
12. (Previously Presented) Process according to Claim 7, wherein the compound of the general formula II or III is selected from 1,1,3,3-tetramethyl-1,3-divinylidisiloxane,

1,3,5,7-tetravinyl-1,3,5,7-tetramethylcyclotetrasiloxane and 1,3,5-trimethyl-1,3,5-trivinylcyclotrisiloxane.

13. (Previously Presented) Process according to Claim 1, wherein the palladium compound is selected from PdX_2 , PdX_4 , M_2PdX_4 , M_2PdX_6 , $(\text{NH}_3)_2\text{PdX}_2$ and $[\text{Pd}(\text{NH}_3)_4]\text{X}_2$, where M is a hydrogen atom, an alkali metal or $\text{NR}^*_4^+$ (R^* = hydrogen, C_{1-4} -alkyl) and X is a halogen or NO_3^- .
14. (Original) Process according to Claim 13, wherein X is chlorine.
15. (Previously Presented) Process according to Claim 1, wherein the reaction is effected in the presence of a solvent or solvent mixture.
16. (Original) Process according to Claim 15, wherein the solvent is selected from water, C_{1-6} -alcohols and C_{2-6} -ethers and mixtures thereof.
17. (Previously Presented) Process according to Claim 1, wherein the base is selected from alkali metal salts, alkaline earth metal salts and ammonium salts (ammonium as NR_4^+ where $\text{R} = \text{H}$ or C_{1-4} -alkyl) of carbonates, hydrogencarbonates and hydroxides.
18. (Previously Presented) Process according to Claim 1, also comprising a purification step.
19. (Previously Presented) Process according to Claim 1, also comprising a concentration step.
20. (Previously Presented) Process according to Claim 1, wherein the reaction of the palladium compound with one or more compounds of the general formula I, II or III is carried out in the presence of one or more ligands other than the compound of the general formula I, II or III.

21. (Previously Presented) Process according to Claim 1, further comprising the reaction of the palladium compound with one or more ligands other than the compound of the general formula I, II or III.
22. (Cancelled)
23. (Previously Presented) Palladium(0) compound obtainable by a process according to Claim 7, wherein the compound of the general formula II is 1,3,5,7-tetravinyl-1,3,5,7-tetramethylcyclotetrasiloxane.